RESEARCH

BMC Pediatrics



Neck circumference cut-off points for detecting overweight and obesity among school children in Northern Cyprus



Ezgi Turkay^{*} and Seray Kabaran

Abstract

Background Neck circumference is one of the anthropometric parameters that is widely used in practical applications, clinical and epidemiological studies in children. It is aimed to determine the neck circumference cut-off points and to use them in the detection of overweight and obesity in children living in Northern Cyprus.

Subjects This cross-sectional study was conducted between October 2019 and January 2020, and covered a sample of 692 children (367 girls and 325 boys) aged 6–10 years attending primary schools in the Northern Cyprus.

Methods Body weight, height, neck circumference, waist circumference, subscapular and triceps skinfold tickness were measured. BF% was calculated with Slaughter equations. World Health Organization BMI cut-off points for age and gender percentiles were used to categorize obesity. BMI, WHtR, NC, body fat were calculated. The Pearson Correlation co-efficient between NC and the other anhtropometric measurements were calculated. Receiver operating characteristics analysis, sensitivity, specificity, PV + ve PV- was used to determine the optimal NC cut-off points for identifying children with overweight and obesity.

Results NC was a statistically significant positive and strong relationship with body weight, BMI, waist circumference and hip circumference (p < 0,005). NC cut-off values to define overweight and obesity were calculated as 26,9 cm in girls (AUC: 0,851, 95% CI: 0,811–0,891, sensitivity 70,50%, specificity 84,65%) and 27,9 cm in boys (AUC: 0,847, 95% CI: 0,805–0,888, sensitivity 76,4%, specificity 79,3%). The ROC curves accurately define overweight and obesity of the whole cohort regardless of age for both sexes of children.

Conclusions The cut-off points for neck circumference were found to be 27,9 cm for boys and 26,9 cm for girls in determining overweight and obesity in children aged 6–10 years. The NC cut-off points obtained in this study can be used to define overweight and obesity in children in epidemiological studies. It is considered to shed light on studies that will examine the relationship between neck circumference and diseases with more people in future studies.

This is a cross-sectional study that asses the correlation between anthropometric adiposity indicators and NC and to establish cut-off points on NC for both sexes at 6-10 years old schoolchildren in Northern Cyprus. The design and conduct of the work was performed by all the authors. The manuscript has been written, read and approved by all the authors. The material has not been published previously, neither whole nor partially and it is also not under consideration for publication elsewhere. Department of Nutrition and Dietetics, Faculty of Health Sciences, Eastern Mediterranean University, T.R. North Cyprus via Mersin 10, Famagusta, Turkey

*Correspondence: Ezgi Turkay ezgii.turkay@gmail.com



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/licenses/by/4.0/. The Creative Commons.org/licenses.org/licenses.org/licenses.org/licenses.org/licen

Keywords Neck circumference, Cut-off points, School children, Overweight and obesity

Introduction

Obesity is a clinical condition that occurs with the increase of adipose tissue in cases where the amount of energy taken into the body is more than the energy expenditure [1]. Obesity has been defined by the World Health Organization (WHO) as "abnormal or excessive fat accumulation in the body to the extent that it impairs health" [2]. The Global Burden of Disease Study has systematically evaluated the prevalence of childhood overweight and obesity since 1980 and has since been shown to have doubled in more than 70 countries worldwide [3]. A total of 107.7 million children (and 603.7 million adults) were classified as obese in 2015, accounting for 23% of the worldwide prevalence of childhood overweight and obesity [4]. It is known that mild obesity is seen between 10-15% and obesity between 1.6-16.0% in children and adolescents in Turkey [5]. In a study conducted in the Turkish Republic of Northern Cyprus, the rates of overweight and obesity in primary school students were found to be 11.3% and 24.6%, respectively [6]. In another study, it was determined that 18.6% of children and adolescents in the 5-19 age group were overweight and 16.2% were obese [7].

Obesity is assuming pandemic proportions and is a major risk factor for cardiometabolic diseases such as diabetes, hypertension, dyslipidemia, and coronary heart disease [8]. Excessive body weight (BW) is the result of complex interactions between genes, dietary intake, physical activity, and the environment. The consequences of excessive childhood BW are a serious public health problem [9]. Childhood obesity requires careful monitoring for early diagnosis and taking precautions for complications that may occur in adulthood. [10].

Anthropometric parameters are the most frequent tools for identifying overweight and obesity because of their practicality. Methods such as body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR) are the most commonly used anthropometric incides [11–14].

BMI is useful to calculate total obesity. WC is used to describe central obesity and abdominal obesity [15, 16]. Because of these anthropometric parameters, which are affected by conditions such as clothing, satiety, and breathing, new strategies are required to find a better scale to measure overweight and obesity, with a particular focus on visceral obesity. [17]. The distribution of adipose tissue in the upper segments of the body, especially with increased visceral adipose tissue is a better predictor of obesity-related complications [18, 19].

In recent years, neck circumference (NC) is a simple, inexpensive, and time-saving anthropometric parameter defined to show central obesity [20–23] and is not affected by conditions such as time of day or season. This measure has optimal results in pediatric assessment and can be used in clinical practice or epidemiological studies [24, 25].

The NC has been recommended as an index of fat distribution in the upper body because subcutaneous fat releases more free fatty acids in the upper part of the body than in the lower part [26]. There is some evidence that a greater NC predicts overweight and obesity [17, 27]. A greater NC, indicator of fat distribution in the upper body, has been shown to be associated with risk of cardiovascular and metabolic disease [28]. It was found that NC was positively correlated with the occurrence of metabolic abnormalities in obese children [29].

In a meta-analysis study results indicated that the NC was a good predictor of elevated blood pressure [30], NC cut-off values were found to be a reliable and easy-to-use tool to determine overweight and obesity in children [31]. Also, another study emphasizes that NC may be considered as an additional significant parameter for monitoring growth and development [32].

To our knowledge, there are no reference data on NC measurements in a large population-based sample of children in Northern Cyprus. This resarch was planed to identify the correlation between NC and anthropometric adiposity indicators and to establish cut-off points on NC for both sexes at 6–10 years old schoolchildren in Northern Cyprus. It is planned to guide the planning of national studies with more people in the future, which also examines its relationship with diseases.

Methods

This is a cross-sectional study carried out among the primary school children, aged 6 to 10 years in Northern Cyprus from October 2019 to January 2020.

The study was approved by the Ethics Committee of Eastern Mediterranean University of Medical Sciences (approval date: 23.05.2019, approval no: 2019/15–10). Approval was obtained from the Turkish Republic of Northern Cyprus, Ministry of National Education, Department of Primary Education (approval date: 05.09.2019, approval no: İÖD.0.00–006-19-E.4541). Consent form was obtained from the parents of the children.

After obtaining written informed consent from parents, a total of 692 students (367 girls and 325 boys) were selected. Northern Cyprus is divided into six districts and twelve sub-districts. There are 32 primary schools at districts areas and 50 primary schools at sub-districts areas. By taking the ratio of the sample number to the total population (n/N), the stratified random sample rate was found according to the districts and the number of schools in the regions to be included in the study was determined. These schools were randomly selected to represent the districts and sub-districts areas. The minimum number of primary school students to be reached is calculated as 546. Exclusion criteria included children with conditions that could affect NC, such as goiter, swelling or cysts in the neck, and cervical spine anomalies, such as cranio-vertebral junction anomalies. Children with growth and developmental delays, who are taking medications that affect the growth and development process, receiving hormone replacement therapy, with amputations and postural deformities were also excluded from the study.

Anthropometric measurements

All anthropometric measurements were conducted by the same trained researcher dietician in accordance with standart procedures.

Body weight

BW was measured without shoes and with light clothing to the nearest 0,1 kg with a calibrated Sinbo SBS-4414 electronic scale [20].

Height

Height was measured with a nonstretchable tape with the subject standing upright, barefoot, and head held in the Frankfurt plane, to the nearest 0,1 cm [33].

Body mass index

The BMI of the subjects was calculated as weight divided by height squared (kg/m²) [34]. World Health Organization BMI cut-offs (WHO) for age and sex percentiles were used to categorize obesity.

Neck circumference

NC was measured to the nearest 0,1 cm with a nonstretchable tape on the midline of the neck among the the cervical backbone and the anterior neck while the subject stood upright, face forward, and shoulders relaxed [35].

Waist circumference

The WC was measured at midpoint between the lowest border of the rib cage and the upper iliac crest to the nearest 0,1 cm [36].

Waist to height ratio

Waist to height ratio (WHtR) was calculated as WC divided by height [37].

Skinfold thickness

The skinfold thickness was measured with the Holtain skinfold caliper to the nearest 0,1 mm while the fingers continued to hold the skinfold. The actual reading was taken approximately 3 s after the caliper tension was released.

Triceps skinfold thickness

Triceps was measured at a marked point in the middle of the posterior surface of the left humerus between the acromion and the olecranon process. The child stood erect, weight was evenly distributed, feet were together, shoulders were relaxed, and arms hung freely at the sides.

Subscapular skinfold thickness

Subscapular circumference was measured while the child stood upright, shoulders were relaxed, and arms hung loosely at the sides. The inferior angle of the left scapula was palpated and a cross was made on the inferior angle of the scapula [38]. Body fat percentage was calculated according to Slaughters' equations [39]. These equations were used because triceps and subcapular skin folds are the most frequently used anthropometric measurements encountered in the literature [40]. All these measurements were taken twice, or three times if differences of more than 2 cm were found, and the average values were calculated.

Slaughter equations [39]: All F: Fat (%) = 1.33(tric + subsc) - 0.013 (tric + subsc)² - 2.5

Prepubertal M: Fat (%) = 1.21 (tric + subsc) - 0.008 (tric + subsc)² - 1.7

All F when (tric + subsc) > 35 mm: Fat (%) = 0.546(tric + subsc) + 9.7

All M when (tric + subsc) > 35 mm: Fat (%) = 0.783(tric + subsc) + 1.7

Statistical analysis

The sample size was calculated as $\alpha = 0.03$, $\beta = 0.05$ assuming independent group means test. The sampling method of the study is stratified random sampling. The formula used for the calculation is given below.

 $n = (Z\alpha/2 + Z\beta)2 * 2 * \sigma 2/d2$

 $Z\alpha/2 =$ normal distribution value of $\alpha/2$,

 $Z\beta/2 =$ normal distribution value of $\beta/2$,

- $\sigma 2 =$ variance of the population and
- d = the difference to be determined.

Descriptive statistics are presented with frequency, percentage, mean, standard deviation, IQR and

minimum-maximum values. The normality assumption was evaluated using Shapiro-Wilks test. Comparison of anthropometric parameters according to gender was compared using the Mann-Whitney U test when the data were not normally distributed and the independent t-test when they were normally distributed. The relationship between the variables was assessed using the Spearman's rank correlation coefficient because of non-normality. Correlation strength was evaluated as low-medium-high [41].

In the multivariate analysis, risk factors for overweight or obesity were examined using binary logistic regression analysis, taking into account possible factors from the univariate analysis. The Hosmer–Lemeshow test was used for model fitting. Receiver operating characteristic analysis (ROC), area under the curve (AUC), sensitivity, specificity, positive predictive value (PV+) and negative predictive value (PV-) was used to determine the optimal NC cut-off points for identifying children with overweight and obesity. Both youden index and sensitivity selectivity were examined to determine the optimal cut-off point in this study. The point with the highest sensitivity selectivity was taken as the cut-off point. The diagnostic test was considered to be "highly accurate if, $0.65 \le AUC \le 1.00^{\circ}$ and "moderately accurate if, $0.50 \le AUC \le 0.65^{\circ}$ [42, 43].

Analyzes were conducted using the Statistical Package for the Social Sciences (SPSS) 23.0 program and p < 0.05 was considered statistically significant.

Results

The total of 692 students, aged 6–10 was assessed anthropometrically. 53% (n=367) of the students participating in the study were girl and 47% (n=325) were boy. The mean BMI was 17,67 kg/m² and 17,85 kg/m², the mean NC was 26,33 cm and 27,49 cm girls and boys respectivelty. Sex-specific mean of other anthropometric measurements are listed in Table 1.

The mean BMI was 15,55 kg/m² and 21,06 kg/m², the mean NC was 25,82 cm and 28,46 cm in normal and overweight/obese children, respectively. There were significant differences between healthy and overweight–obese children with respect to anthropometric parameters (Table 2).

Odds ratios (ORs) were also calculated to determine the strength of association between NC and overweight/ obesity. 1 unit increase in NC poses 1,519 times higher risk of being overweight and obese (Table 3).

Tab	le	1 (Com	parison	of a	nthro	oometric	parameters	by gende	er

Anthropometric SD Median IQR Min n Mean Max р parameters BW (kg) Boy 325 30,98 9,12 29 10,2 16,4 67 Girl 367 30,11 9,22 27,9 10,8 15,4 74,3 0,106 Height (cm) 103 Boy 325 130,74 9.87 130 14.5 156 Girl 367 129,41 9,65 129,5 14 104 158 0,075 $BMI (kg/m^2)$ Boy 325 1785 3.58 17.01 3 68 10.34 37.31 0,358 Girl 367 17,67 3,48 16,73 4,05 12,3 33,14 WC (cm) 9,84 Boy 325 60,6 57,5 11 44 104 Girl 367 58,79 8,86 56 12 45 94 0,009 NC (cm) 325 27,49 2,27 27 2,5 23 36 Bov 2,19 20,5 Girl 367 26,33 26 2,7 35,5 < 0.0001 WHtR Boy 306 0,46 0.06 0.45 0,07 0,36 0.78 Girl 339 0,45 0,05 0,06 0,35 0,7 0,196 0.44 Body fat (%) 306 7,59 9,56 Boy 18.31 16.37 7.69 43.67 Girl 350 1972 5 98 18,9 8,01 9,32 45,01 < 0,0001

Mann-Whitney U test was used for all parameters. IQR Interquartile range: Q3-Q1

BW Body weight, BMI Body mass index, WC Waist circumference, NC Neck circumference, WHtR Wasit-to-height raito

	BMI for age	n	Mean	Std. Dev	Min	Max	p
BW (kg)	N*	415	25,87	4,86	15,4	42,3	< 0,0001
	O*	277	37,5	9,69	20,4	74,3	
Height (cm)	N*	415	128,45	9,51	104	157	< 0,0001
	O*	277	132,41	9,7	103	158	
BMI (kg/m ²)	N*	415	15,55	1,35	10,34	18,82	< 0,0001
	O*	277	21,06	3,19	17	37,31	
NC (cm)	N*	415	25,82	1,52	21	31,5	< 0,0001
	O*	277	28,46	2,35	20,5	36	
WC (cm)	N*	415	54,61	4,88	44	85	< 0,0001
	O*	277	67,16	9,46	50	104	
WHtR	N*	415	0,43	0,03	0,35	0,6	< 0,0001
	O*	277	0,51	0,05	0,39	0,78	
Body fat (%)	N [*]	409	15,54	4,11	7,69	38,4	< 0,0001
	O*	247	24,9	6,37	10,58	45,01	

Table 2 Standard deviation, median, min-max and p values of anthropometric parameters according to overweight/obesity distribution

İndependent t test used for all parameters

BW Body weight, BMI Body mass index, WC Waist circumference, NC Neck circumference, WHtR Wasit-to-height raito

Table 3	Indepen	dent risk ⁻	factor for	being o	verweight a	and o	bese
---------	---------	------------------------	------------	---------	-------------	-------	------

Risk Factor	OR (95% CI)	<i>p</i> value	
NC	1,519 (1,291–1,788)	< 0,0001	

 χ^2 = 13,44 p = 0,098 (Hosmer and Lemeshow Test), Nagelkerke R^2 = 0,637 NC Neck circumference, OR Odds ratio Table 4 shows the correlations between anthropometric variables. NC was a statistically significant positive and strong relationship with body fat, BMI and WC. The relationship between anthropometric parameters according to gender is also presented separately in Table 4. All

Table 4 Correlation co-efficient between neck circumference and adiposity anthropometric indicators in children

Sex	Anthropometric parameters	NC	BMI	WC	WHtR	Body fat
Воу	NC (cm)	-	,760 ^a	,856ª	,650 ^a	,724 ^a
	BMI (kg/m ²)		-	,811 ^a	,814 ^a	,789 ^a
	WC (cm)			-	,860 ^a	,850 ^a
	WHtR				-	,837 ^a
	Body Fat (%)					-
Girl	NC (cm)	-	,700 ^a	,768 ^a	,552 ^a	,623 ^a
	BMI (kg/m ²)		-	,855ª	,800 ^a	,826 ^a
	WC (cm)			-	,834 ^a	,818 ^a
	WHtR				-	,744 ^a
	Body Fat (%)					-
	NC (cm)	-	,710 ^a	,801ª	,598 ^a	,612ª
Total	BMI (kg/m ²)		-	,843 ^a	,820 ^a	,805 ^a
	WC (cm)			-	,850 ^a	,814 ^a
	WHtR				-	,779 ^a
	Body Fat (%)					-

Spearman's rank correlation coefficient was used

NC Neck circumference, BMI Body mass index, WC Waist circumference, WHtR Waist-to-height raito

^a Correlation is significant at the 0.01 level (2-tailed)

parameters have a strong statistically significant positive correlation with each other.

Based on ROC analysis, sensitivities, specificities, and cut-off values for NC for children are presented in Table 5. NC cut-off values to define overweight and obesity were calculated as 26,9 cm in girls (sensitivity 70,50%, specificity 84,65%) and 27,9 cm in boys (sensitivity 76,4%, specificity 79,3%). Also, NC was calculated age and sexspecific for detecting overweight and obesity. NC cut-off values for overweight and obesity increased from 26,0 to 28,8 cm for boys (95% CI: 0,794–0,931) 25.5 to 27,8 cm for girls (95% CI: 0,747–0,934) and 26,9 cm in girls (95% CI: 0,811–0,891) 27,9 cm in boys (95% CI: 0,805–0,888) between 6 and 10 years.

The ROC curves accurately define overweight and obesity of the whole cohort regardless of age for both sexes of children (Fig. 1).

Discussion

The purpose of this study was to investigate whether NC is an alternative method for detecting overweight and obesity in the pediatric population. The cut-off points for NC were found to be 27,9 cm for boys and 26,9 cm for girls in determining overweight and obesity in children aged 6–10 years. Also, we found positive strong relationship with NC and WC, BW and BMI. There is no statistical difference between boys and girls when evaluating BW, height, BMI and WHtR parameters. WC, NC are higher in boys while body fat percentage is higher in girls than boys. Similar to our study, no statistical difference was found between anthropometric parameters except NC and WC in children aged 6–11 years in Western Mexico population [44].

When the mean values of NC according to BW are evaluated it was found in a study that the mean NC values of children with normal BMI were 29.0 and 28.2 cm in boys and girls, respectively, whereas this value was 32.7 and 31.1 cm in overweight and obese children, respectively [45]. In another study, the NC was 31.2 cm in overweight/obese boys and 29.9 cm in girls at 6-10 years of age [46]. As expected, these results showed the same results with our study that overweight/obese children have a larger NC. When WC, WHtR and body fat were examined, the mean value WC, WHtR and body fat ratio of boys with normal BMI were found to be 54.9 cm, 043 and 16.6%, respectively, while these values were 66.8 cm, 0,50 and 25.2% in obese children. These values were found to be 54.9 cm, 0,43 and 18.8% in girls with normal BMI, and 69.1 cm, 0,52 and 31.6% in obese children, respectively [47]. Our study also found similar results with this consequences.

In this study, a strong positive correlation was found between the NC and other anthropometric parameters, the highest correlation with WC, BW and BMI, respectivelly. These results are in agreement with the results of other studies in a similar age group [48, 49]. In another studies, it was found that NC was most correlated with height, BW and WC [27, 50]. In a study examining the relationship between obesity status and NC, BMI and WC were found to be the anthropometric parameters most correlated with neck circumference in both boys and girls [51].

Reported in a study that NC positively correlated with BMI and WC [18]. Similarly, the results of Wang's study showed a significant positive correlation between NC, WC and BMI in Chinese Yi adolescents [52]. These findings are consistent with our study and it shows that

Sex	Age	AUC	p value	cut-off point (cm)	95% CI	$\mathbf{PV}+$	PV-	Sensitivity(%)	Specificity(%)
Воу	6	0,897	< 0,001	>26	0,794–0,959	94,7	90,9	81,82	97,56
	7	0,829	< 0,001	>26,5	0,724–0,906	69,0	84,8	85,29	68,29
	8	0,880	< 0,0001	>27,4	0,775–0,948	87,0	80,5	71,43	91,67
	9	0,943	< 0,001	> 28,5	0,854–0,986	95,0	86,0	76,0	97,37
	10	0,853	< 0,001	> 28,8	0,738–0,931	82,1	81,2	79,31	83,87
Total	6-10	0,847	< 0,001	> 27,9	0,805–0,888	94,7	90,9	76,4	79,3
Girl	6	0,856	< 0,0001	> 25,5	0,747-0,931	62,5	92,7	83,33	80,85
	7	0,876	< 0,001	> 25,7	0,791–0,936	65,0	90,2	83,87	76,67
	8	0,910	< 0,001	> 26,7	0,822–0,963	90,6	84,1	80,56	92,50
	9	0,842	< 0,001	> 26,7	0,741–0,915	70,6	88,4	82,76	79,17
	10	0,855	< 0,001	>27,8	0,738–0,934	74,1	83,9	80,0	78,79
Total	6-10	0,851	< 0,001	> 26,9	0,811–0,891	73,7	82,5	70,50	84,65

Table 5 The values of the area under the curve (AUC), the cut-off value, sensitivity, and specificity of NC in detecting overweight/ obesity in boys and girls according to age

AUC Area Under the Curve, CI Confidence interval, PV + Positive predictive value, PV- Negative predictive value





Boys









b









100-Specificity









I

Fig. 1 Receiver operating characteristic (ROC) curve for determining the optimal NC cut-off values for identifying overweight and obesity in boys and girls. (For boys; a: 6–10 years, b: 6 years, c: 7 years, d: 8 years, e: 9 years, f: 10 years). (For girls; g: 6–10 years, h: 6 years, i: 7 years, j: 8 years, k: 9 years, f: 10 years). (For girls; g: 6–10 years, h: 6 years, i: 7 years, j: 8 years, k: 9 years, f: 10 years).

in circumstances where these methods are not feasible, measurement of NC may be an alternative.

Previous studies suggest that using the NC, which is easy to practise and inexpensive, is a more useful parameter that is not influenced by hunger or satiety or by respiratory movements and provides more stable results for indicating fat accumulation on the upper body. Therefore, it is highly recommended to use it to detect body fat accumulation and associated risk factors [53, 54].

The AUC results show that NC values have highly accurate in detecting overweight and obesity in children of all ages and genders. Moreover, the cut-off values of NC for detecting overweight and obesity in children of different ages were 26-28,8 cm in boys and 25,5-27,8 cm in girls. Different epidomiological studies showed different NC cut-off values in children [31, 43, 55]. In Han children aged 7-12 years, NC cut-off points were 27,4-31,3 cm in boys and 26,3-31,4 cm in girls [55]. In a study conducted in Turkey studied Turkish children aged 6-10 years and found that NC cut-off scores increased from 28,0 to 31,5 for boys and 27,0-30 for girls [32]. In another study in Mexico, NC cut-off scores identifying increased central adiposity at age 6–10 years ranged from 27,5 to 30,2 cm for boys and from 25,7 to 29,2 cm for girls [44]. The differences between the results may be due to differences in age, sociodemographic factors, and ethnicity.

The NC is used not only to determine overweight and obesity, but also to predict diseases that affect adulthood, such as the metabolic syndrome [55, 56]. Studies on NC have shown that there is an association between NC and central obesity and cardiometabolic disorders [44, 30]. A systematic review concluded that NC has predictive value for the diagnosis of some cardiometabolic risk factors, such as a positive association between NC and glycemic indices and lipid profile in adult and prepubertal populations [57]. Demonstrated in the study that NC correlated with blood pressure, triglycerides, and markers of insulin resistance in both gender [58].

In addition to the studies mentioned above, that NC may be an indicator of overweight and obesity in children, NC was found to be lower than WC as a screening tool in a study, and it was stated that more studies are needed to be an indicator of adiposity in children [27]. In a study examining the relationship between NC and cardiometabolic risk factors, it was not found to be associated with risk factors independent of BMI and was evaluated as a poor classifier of cardiometabolic risk factors in children [59]. It is necessary to support these findings with comprehensive studies that will examine the relationship between NC and disease in future.

Strength and limits

The sample size based on the population of the Turkish Republic of Northern Cyprus is a strength of this study and makes the results generalizable. The application of the anthropometric measurements in the research by a single person strengthens the study in terms of ensuring the criteria and standardization of the measurements. The utility of NC as a screening tool for detecting obesity is that it has been evaluated with BMI as the standard criterion, rather than dualenergy X-ray absorptiometry and methods that analyze body composition. One of the limitations of the study was that other anthropometric parameters such as WC were ignored and only NC cut-off values were focused on. Another limitation is that due to the age of the children, the exact dates of birth could not be taken and they could not be calculated as months-years because their age was questioned, not the date of birth. Therefore, to compare BMI percentiles for age from anthropometric measurements of adolescents with those recommended by WHO; Adolescents were considered to be in the sixth month of their current age. One of the limitations of the study is that the body fat ratios of the children were calculated only with the equations.

It is recommended to examine the relationship of neck circumference, which is very practical to use in epidemiological studies, to larger masses, not only in the determination of overweight and obesity, but especially with nutrition-related diseases.

Conclusions

It is recommended to use NC in the determination of overweight and obesity in children and in epidemiological studies because it is a practical, simple, and inexpensive method that is not influenced by hunger-satiety situations. This is the first study to use the NC cut-off point in the determination of overweight and obesity in school-age children in Northern Cyprus. NC cut-off values to define overweight and obesity were calculated as 26,9 cm in girls (sensitivity 70,50%, specificity 84,65%) and 27,9 cm in boys (sensitivity 76,4%, specificity 79,3%). The cut-off values acquired in this study could be used in populations similar to our characteristics properties and lead to creating reference values for wider age groups in Northern Cyprus.

Abbreviations

3W	Body weight
ЗMI	Body mass index
NC	Waist circumference
WHR	Waist-to-hip ratio
NC	Neck circumference
NHO	World Health Organization

- WHtR Waist-to-height ratio ROC Receiver operating characteristics SPSS Statistical Package for the Social Sciences Area Under the Curve AUC CL Confidence interval PV⁺ Positive predictive value PV' Negative predictive value N Normal weight Overweight/obese 0
- OR Odds ratio

Acknowledgements

We are grateful to the participants for their willingness to participate in this study.

Conflict of interest

The authors declare that there is no conflict of interest.

Authors' contributions

Conception: Ezgi Türkay, Seray Kabaran. Design: Ezgi Türkay, Seray Kabaran. Control/Supervision: Seray Kabaran. Data collection and/or processing: Ezgi Türkay. Analysis and/or interpretation: Ezgi Türkay, Seray Kabaran. Literature review: Ezgi Türkay, Seray Kabaran. Writing the manuscript: Ezgi Türkay. Table and figure preparation: Ezgi Türkay. Critical review: Seray Kabaran. The manuscript has been written, read, and approved by all the authors. All authors reviewed the manuscript.

Funding

This research did not receive any financial support and any other funding.

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The study was approved by the Ethics Committee of Eastern Mediterranean University of Medical Sciences (approval date: 23.05.2019, approval no: 2019/15–10). Approval was obtained from the Turkish Republic of Northern Cyprus, Ministry of National Education, Department of Primary Education (approval date: 05.09.2019, approval no: İÖD.0.00–006-19-E.4541). Informed Consent form was obtained from the parents of the children. All methods were performed in accordance with the relevant guidelines and regulations by including a statement.

Consent for publication

This study does not contain identifying images or other personal or clinical details of participants that compromise anonymity.

Competing interests

There are no financial and non-financial competitive interests. The datasets generated and/or analysed during the current study are not publicly available due just to make sure it's used for scientific purposes only but are available from the corresponding author on reasonable request.

Received: 12 November 2021 Accepted: 16 September 2022 Published online: 14 October 2022

References

- Blüher M. Obesity: global epidemiology and pathogenesis. Nat Rev Endocrinol. 2019;15(5):288–98.
- World Health Organization. World health statistics 2016: monitoring health for the SDGs sustainable development goals. Geneva: World Health Organization; 2016.

- GBD 2015 Obesity Collaborators. Health effects of overweight and obesity in 195 countries over 25 years. New Engl J Med. 2017;377(1):13–27.
- Bakanlığı TS. Türkiye Beslenme ve Sağlık Araştırması 2010: Beslenme durumu ve alışkanlıklarının değerlendirilmesi sonuç raporu. Ankara, Sağlık Bakanlığı Sağlık Araştırmaları Genel Müdürlüğü, 2014; 27
- Elmas C. Gazimağusa'daki Devlet İlkokullarında 4. ve 5. Sınıf Öğretmenlerine Verilen Beslenme Eğitiminin Öğrencilere Yansımasının İzlenmesi (Master's thesis, Eastern Mediterranean University EMU-Doğu Akdeniz Üniversitesi (DAÜ), 2017.
- Kabaran S, Gezer C. Kuzey Kıbrıs Türk Cumhuriyeti'ndeki çocuk ve adolesanlarda Akdeniz diyetine uyum ile obezitenin belirlenmesi. Türkiye Çocuk Hastalıkları Dergisi. 2013;7(1):11–20.
- Joshipura K, Muñoz-Torres F, Vergara J, Palacios C, Pérez CM. Neck circumference may be a better alternative to standard anthropometric measures. J Diabetes Res. 2016;2016:1–8.
- Kroll C, Mastroeni SS, Czarnobay SA, Ekwaru JP, Veugelers PJ, Mastroeni MF. The accuracy of neck circumference for assessing overweight and obesity: a systematic review and meta-analysis. Ann Hum Biol. 2017;44(8):667–77.
- Magalhães EIDS, Sant'Ana LFDR, Priore SE, Franceschini SDCC. Waist circumference, waist/height ratio, and neck circumference as parameters of central obesity assessment in children. Revista Paulista de Pediatria. 2014;32(3):273–81.
- Zvonar M, Štefan L, Kasović M. Percentile curves for body-mass index, waist circumference, waist-to-height ratio and waist-to-height ratio (Exp) in Croatian adolescents. Int J Environ Res Public Health. 2019;16(11):1920.
- 12. Aggarwal B, Jain V. Obesity in children: definition, etiology and approach. Indian J Pediatr. 2018;85(6):463–71.
- Asif M, Aslam M, Ullah K, Qasim M, Afzal K, Abbas A, et al. Diagnostic performance and appropriate cut-offs of different anthropometric indicators for detecting children with overweight and obesity. BioMed Res Int. 2021;2021:1-11.
- Alves Junior CAS, Mocellin MC, Gonçalves ECA, Silva DAS, Trindade EBSM. Anthropometric indicators as body fat discriminators in children and adolescents: a sys- tematic review and meta-analysis. Adv Nutr. 2017;8(5):718–27.
- Adom T, Kengne AP, De Villiers A, Boatin R, Puoane T. Diagnostic accuracy of body mass index in defining childhood obesity: analysis of crosssectional data from Ghanaian children. Int J Environ Res Public Health. 2020;17(1):36.
- Martin-Calvo N, Moreno-Galarraga L, Martinez-Gonzalez MA. Association between body mass index, waist-to-height ratio and adiposity in children: a systematic review and meta-analysis. Nutrients. 2016;8(8):512.
- 17. Taheri M, Kajbaf TZ, Taheri MR, Aminzadeh M. Neck circumference as a useful marker for screening overweight and obesity in children and adolescents. Oman Med J. 2016;31(3):170.
- Jindal M, Masand R, Nimawat AK, Natani BS, Verma CR. Neck circumference and waist circumference as a tool for evaluating obesity. Indian J Child Health. 2020;7(4):159–63.
- Taghizadeh S, Farhangi MA. Neck Circumference as a Potent Anthropometric Predictor of Cardiovascular Diseases and Metabolic Syndrome: A Review Study. J Fasting Health. 2019;7(2):70–5.
- Mucelin E, Traebert J, Zaidan MA, Piovezan AP, Nunes RD, Traebert E. Accuracy of neck circumference for diagnosing overweight in six and seven-year-old children. Jornal de Pediatria. 2021;2021(97):559–63.
- Malini M, Kalpana S, Lakshmi S. Neck circumference in children and adolescents-an emerging tool for screening central obesity. Indian J Child Health. 2022;9(1):11–5.
- 22. Folmann AG, Wolf VLW, Roman EP, Guerra-Júnior G. Neck circumference and excess weight: proposal of cutoff points for Brazilian adolescents. Jornal de Pediatria. 2021;97:191–6.
- 23. Sreelatha PR, Chinchilu RV. Neck circumference-A simple and valid screening tool for obesity in school children. J Surg Med. 2021;5(12):1184–7.
- 24. Kumar NV, Ismail MH, Mahesha P, Girish M, Tripathy M. Neck circumference and cardio-metabolic syndrome. J Clin Diagn Res. 2014;8(7):MC23.
- González-Cortés CA, Téran-García M, Luevano-Contreras C, Portales-Pérez DP, Vargas-Morales JM, Cubillas-Tejeda AC, et al. Neck Circumference and

Its Association with Cardiometabolic Risk Factors in Pediatric Population. Medicina. 2019;55(5):183.

- Maskey M, Gupta KD, Ahmed M. Neck Circumference, a Novel Predictor of Overweight/Obesity in School Children in Pokhara. Med Phoenix. 2020;5(1):32–9.
- 27. Kelishadi R, Djalalinia S, Motlagh ME, Rahimi A, Bahreynian M, Arefirad T, et al. Association of neck circumference with general and abdominal obesity in children and adolescents: the weight disorders survey of the CASPIAN-IV study. BMJ Open. 2016;6(9):e011794.
- Katz SL, Vaccani JP, Clarke J, Hoey L, Colley RC, Barrowman NJ. Creation of a reference dataset of neck sizes in children: standardizing a potential new tool for prediction of obesity-associated diseases? BMC Pediatr. 2014;14(1):159.
- Kurtoglu S, Hatipoglu N, Mazicioglu MM, Kondolot M. Neck circumference as a novel parameter to determine metabolic risk factors in obese children. Eur J Clin Invest. 2012;42(6):623–30.
- Moradi S, Mohammadi H, Javaheri A, Ghavami A, Rouhani MH. Association between neck circumference and blood pressure: a systematic review and meta-analysis of observational studies. Horm Metab Res. 2019;51(8):495–502.
- Hatipoglu N, Mazicioglu MM, Kurtoglu S, Kendirci M. Neck circumference: an additional tool of screening overweight and obesity in childhood. Eur J Pediatr. 2010;169(6):733–9.
- Mazicioglu MM, Kurtoglu S, Ozturk A, Hatipoglu N, Cicek B, Ustunbas HB. Percentiles and mean values for neck circumference in Turkish children aged 6–18 years. Acta Paediatr. 2010;99(12):1847–53.
- Xiong KY, He H, Zhang YM, Ni GX. Analyses of body composition charts among younger and older Chinese children and adolescents aged 5 to 18 years. BMC Public Health. 2012;12(1):1–9.
- Sarria A, Moreno LA, Garcí-Llop LA, Fleta J, Morellon MP, Bueno M. Body mass index, triceps skinfold and waist circumference in screening for adiposity in male children and adolescents. Acta Paediatr. 2001;90(4):387–92.
- 35. Stabe C, Vasques ACJ, Lima MMO, Tambascia MA, Pareja JC, Yamanaka A, et al. Neck circumference as a simple tool for identifying the metabolic syndrome and insulin resistance: results from the Brazilian Metabolic Syndrome Study. Clin Endocrinol. 2013;78(6):874–81.
- Bacopoulou F, Efthymiou V, Landis G, Rentoumis A, Chrousos GP. Waist circumference, waist-to-hip ratio and waist-to-height ratio reference percentiles for abdominal obesity among Greek adolescents. BMC Pediatr. 2015;15(1):1–9.
- Kuciene R, Dulskiene V. Associations between body mass index, waist circumference, waist-to-height ratio, and high blood pressure among adolescents: a cross-sectional study. Sci Rep. 2019;9(1):1–11.
- Rodriguez G, Moreno LA, Blay MG, Blay VA, Fleta J, Sarria A, et al. Body fat measurement in adolescents: comparison of skinfold thickness equations with dual-energy X-ray absorptiometry. Eur J Clin Nutr. 2005;59(10):1158–61.
- Slaughter MH, Lohman TG, Boileau RA, Horswill CA, Stillman RJ, van Loan MD, et al. Skinfold equations for estimation of body fatness in children and youths. Hum Biol. 1988;60:709–23.
- Cerqueira MS, Amorim PR, Encarnação IG, Rezende LM, Almeida PH, Silva AM, et al. Equations based on anthropometric measurements for adipose tissue, body fat, or body density prediction in children and adolescents: a scoping review. Eat Weight Disord Stud Anorexia, Bulimia Obesity. 2022; 1–18.
- 41. Schober P, Boer C, Schwarte LA. Correlation coefficients: appropriate use and interpretation. Anesth Analg. 2018;126(5):1763–8.
- Perkins NJ, Schisterman EF. The inconsistency of "optimal" cutpoints obtained using two criteria based on the receiver operating characteristic curve. Am J Epidemiol. 2006;163:670–5 (Epub 2006 Jan 12).
- Hanley JA. The robustness of the "binormal" assumptions used in fitting ROC curves. Med Deci Mak. 1988;8:197–203.
- Valencia-Sosa E, Chávez-Palencia C, Romero-Velarde E, Larrosa-Haro A, Vásquez-Garibay EM, et al. Neck circumference as an indicator of elevated central adiposity in children. Public Health Nutr. 2019;22(10):1755–61.
- Patnaik L, Pattnaik S, Rao EV, Sahu T. Validating neck circumference and waist circumference as anthropometric measures of overweight/obesity in adolescents. Indian Pediatr. 2017;54(5):377–80.
- 46. Nafiu OO, Burke C, Lee J, Voepel-Lewis T, Malviya S, Tremper KK. Neck circumference as a screening measure for identifying children with high body mass index. Pediatrics. 2010;126(2):e306–10.

- Coutinho CA, Longui CA, Monte O, Conde W, Kochi C. Measurement of neck circumference and its correlation with body composition in a sample of students in São Paulo. Brazil Hormone Res Paediatr. 2014;82(3):179–86.
- Asif M, Aslam M, Tariq Ismail SA, Rahman A. Neck Circumference as Screening Measure for Identifying Children with Central Obesity, A study from Pakistan. Pak Pediatr J. 2020;44(3):277–83.
- Asif M, Aslam M, Wyszyńska J, Altaf S, Ahmad S. Diagnostic performance of neck circumference and cut-off values for identifying overweight and obese pakistani children: a receiver operating characteristic analysis. J Clin Res Pediatr Endocrinol. 2020;12(4):366.
- Katamba G, Agaba DC, Muzaale F, Namaganda A, Namayanja R, Musasizi A, et al. The Use of Neck Circumference as Marker of Overweight and Obesity Among Adolescents Aged 12–19 Years. A Case of Southwestern Uganda Secondary School Adolescents. 2020.
- Nafiu OO, Zepeda A, Curcio C, Prasad Y. Association of neck circumference and obesity status with elevated blood pressure in children. J Hum Hypertens. 2014;28(4):263–8.
- Wang H. Analyzing Neck Circumference as a Tool for Evaluating Overweight and Obesity in Chinese Adolescents. J Healthcare Eng. 2021;2021(2021):1–6.
- Ma C, Wang R, Liu Y, Lu Q, Liu X, Yin F. Diagnostic performance of neck circumference to identify overweight and obesity as defined by body mass index in children and adolescents: systematic review and meta-analysis. Ann Hum Biol. 2017;44(3):223–9.
- Atwa H, Fiala L, Handoka NM. Neck circumference as an additional tool for detecting children with high body mass index. J Am Sci. 2012;8(10):442–6.
- Lou DH, Yin FZ, Wang R, Ma CM, Liu XL, Lu Q. Neck circumference is an accurate and simple index for evaluating overweight and obesity in Han children. Ann Hum Biol. 2012;39(2):161–5.
- Ataie-Jafari A, Namazi N, Djalalinia S, Chaghamirzayi P, Abdar ME, Zadehe SS, et al. Neck circumference and its association with cardiometabolic risk factors: a systematic review and meta-analysis. Diabetol Metabol Syndr. 2018;10(1):1–34.
- Castro-Piñero J, Delgado-Alfonso A, Gracia-Marco L, Gómez-Martínez S, Esteban-Cornejo I, Veiga OL, et al. Neck circumference and clustered cardiovascular risk factors in children and adolescents: cross-sectional study. BMJ Open. 2017;7(9):e016048.
- Silva CD, Zambon MP, Vasques ACJ, Rodrigues AMDB, Camilo DF, Antonio MÂR, et al. Neck circumference as a new anthropometric indicator for prediction of insulin resistance and components of metabolic syndrome in adolescents: Brazilian Metabolic Syndrome Study. Revista Paulista de Pediatria. 2014;32:221–9.
- Shirley MK, Alves Pereira-Freire J, de Macêdo Gonçalves Frota K, Oliveira Lemos J, Wells JC, Arnaud Rosal Lopes Rodrigues L, et al. Evaluation of neck circumference as a predictor of elevated cardiometabolic risk outcomes in 5–8-year-old Brazilian children. Child Adolesc Obesity, 2020; 3(1), 1–19.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

